

Comment on “Solitonlike Solutions of the Grad-Shafranov Equation”

G. N. Throumoulopoulos^{*1}, K. Hizanidis^{#2}, H. Tasso^{†3}

**University of Ioannina,*

Association Euratom-Hellenic Republic,

Department of Physics, GR 451 10 Ioannina, Greece.

#National Technical University of Athens,

School of Electrical and Computer Engineering,

9 Iroon Polytechniou Str., Zografou Campus,

157 73 Athens, Greece

† Max-Planck-Institut für Plasmaphysik,

Euratom association,

D-85748 Garching bei München, Germany

Abstract

In the above entitled recent publication by Giovanni Lapenta [Phys. Rev. Lett. **90**, 135005 (2003)] it is claimed construction of a new class of soliton-like solutions for the Grad-Shafranov equation in plane geometry. It is proved here that, because of the mathematically erroneous choice $\nabla p = |\Psi|^2 \Psi \nabla \Psi$ for an analytic continuation of the poloidal magnetic flux-function Ψ in the complex plane (p is the pressure), the cubic Schrödinger equation considered by the author is irrelevant to the equilibrium problem and the Grad-Shafranov equation.

¹gthroum@artemis1.physics.uoi.gr

²kyriakos@central.ntua.gr

³het@ipp.mpg.de

In a recent publication [1], the author claims derivation of a new class of solitonlike solutions for the Grad-Shafranov equation in plane geometry. The equilibrium equations considered are (Eqs. (5) [1])

$$\begin{aligned} (\nabla\Psi \times \nabla B_z) \cdot \hat{\mathbf{z}} &= 0 \\ \nabla p + \nabla^2\Psi \nabla\Psi + B_z \nabla B_z &= 0, \end{aligned} \quad (1)$$

where $p(\Psi)$ and $B_z(\Psi)$ are the pressure and the z -component of the magnetic field, respectively. The following forms of the free functions p and B_z are then chosen (Eq. (13) [1])

$$B_z \nabla B_z = \alpha_0^2 \Psi \nabla \Psi \quad (2)$$

$$\nabla p = \alpha_0^2 |\Psi|^2 \Psi \nabla \Psi, \quad (3)$$

and (1) is extended in the complex plane, thus leading to the cubic Schrödinger equation (Eq. (14) [1])

$$\frac{\partial^2 \Psi}{\partial x^2} + \frac{\partial^2 \Psi}{\partial y^2} = -\alpha_0^2 (1 + |\Psi|^2) \Psi. \quad (4)$$

A solitonlike solution of (4) is (Eq. (15) [1])

$$\Psi(x, y) = \Psi_p \operatorname{sech}(x/L) e^{-j(\alpha_0 + 1/2\alpha_0 y_0^2)y}. \quad (5)$$

For a complex function Ψ , however, the rhs of (3) becomes a non holomorphic function, i.e. owing to the fact that $|\Psi|$ does not have derivative, the term $\alpha_0^2 |\Psi|^2 \Psi \nabla \Psi$ can not be a function gradient as the lhs of (3) requires. An explicit proof follows. Taking the curl of (3) yields $\nabla|\Psi| \times \nabla\Psi = 0$, implying that Ψ depends only on $|\Psi|$:

$$\Psi = f(|\Psi|). \quad (6)$$

In order that the complex function Ψ be analytic, on account of (6) and the Cauchy-Riemann conditions leads to $|\Psi| = \text{constant}$ and therefore $\Psi = \text{constant}$. Also, even without requiring analyticity of Ψ , by only considering its polar form, $\Psi = |\Psi| \exp(j\Theta(x, y))$, Eq. (6) implies that

$$\Theta = \Theta(|\Psi|). \quad (7)$$

Solution (5), however, is inconsistent with (7) (otherwise it should hold that $x = x(y)$). Therefore, (4) is irrelevant to the Grad-Shafranov equation; as a matter of fact the real part of (5),

$$u(x, y) = \Psi_p \operatorname{sech}(x/L) \cos \left[(\alpha_0 + 1/2\alpha_0 y_0^2) y \right],$$

does not satisfy the respective Grad-Shafranov equation

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = -\alpha_0^2(1 + u^2)u. \quad (8)$$

A mathematically legitimate choice for ∇p , instead of (3), could be

$$\nabla p = \alpha_0^2 \Psi^3 \nabla \Psi. \quad (9)$$

This leads to an equation of the form (8) for Ψ . Solving this equation in the complex plane, however, is a task not easier than that for real Ψ .

In conclusion, because of the mathematically erroneous choice (3) for an analytic continuation of Ψ , the cubic Schroedinger equation (4) considered by the author is irrelevant to the equilibrium problem (1) and to the Grad-Shafranov equation. Despite this unlucky situation we consider the idea of the author as appealing, and hope that it will be successful in the future if used in an appropriate setting.

Acknowledgements

Part of this work was conducted during a visit of one of the authors (G.N.T.) to the Max-Planck Institut für Plasmaphysik, Garching. The hospitality of that Institute is greatly appreciated.

The present work was performed under the Contract of Association ERB 5005 CT 99 0100 between the European Atomic Energy Community and the Hellenic Republic.

References

- [1] Giovanni Lapenta, Phys. Rev. Lett, **90**, 135005 (2003).